arXiv:1112.2949v2 [math.AC] 28 Mar 2012

FUNDAMENTAL INVARIANTS FOR THE ACTION OF $SL_3(\mathbb{C}) \times SL_3(\mathbb{C}) \times SL_3(\mathbb{C})$ ON $3 \times 3 \times 3$ ARRAYS

MURRAY R. BREMNER AND JIAXIONG HU

Abstract. We determine the three fundamental invariants in the entries of a $3 \times 3 \times 3$ array over $\mathbb{C}$ as explicit polynomials in the 27 variables x_{ijk} for $1 \le i, j, k \le 3$. By the work of Vinberg on θ-groups, it is known that these homogeneous polynomials have degrees 6, 9 and 12; they freely generate the algebra of invariants for the Lie group $SL_3(\mathbb{C}) \times SL_3(\mathbb{C}) \times SL_3(\mathbb{C})$ acting irreducibly on its natural representation $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$. These generators have respectively 1152, 9216 and 209061 terms; we find compact expressions in terms of the orbits of the finite group $(S_3 \times S_3 \times S_3) \rtimes S_3$ acting on monomials of weight zero for the action of the Lie algebra $\mathfrak{sl}_3(\mathbb{C}) \oplus \mathfrak{sl}_3(\mathbb{C}) \oplus \mathfrak{sl}_3(\mathbb{C})$.

1. Introduction

We consider $3 \times 3 \times 3$ arrays $X = (x_{ijk})$ where $x_{ijk} \in \mathbb{C}$ for $1 \le i, j, k \le 3$. We are concerned with homogeneous polynomials in the entries x_{ijk} of these arrays which are invariant under the action of the Lie group $SL_3(\mathbb{C}) \times SL_3(\mathbb{C}) \times SL_3(\mathbb{C})$ acting by simultaneous changes of basis along the three directions.

Following Bremner et al. [2, 3], we use a computational approach, based on the representation theory of the special linear Lie algebra $\mathfrak{sl}_3(\mathbb{C})$, to express these invariant polynomials as elements in the nullspace of a large integer matrix, and then to determine explicitly the generators of the algebra of invariants. Apart from their intrinsic interest, these generators can in principle be used to determine an explicit form for the hyperdeterminant of a $3 \times 3 \times 3$ array in the sense of Gelfand et al. [8]; see the open problem at the end of this paper.

Vinberg [16] generalized the notion of Weyl group from Lie groups to θ-groups; a θ-group is the group of fixed points of an automorphism of finite order of a complex semisimple Lie group. He showed that the corresponding Weyl groups are generated by complex reflections, and obtained the corollary that the algebra of invariants is a polynomial algebra: it is freely generated by a finite set of fundamental invariants. In particular, he embedded the Lie algebra $\mathfrak{sl}_{333}(\mathbb{C}) = \mathfrak{sl}_3(\mathbb{C}) \oplus \mathfrak{sl}_3(\mathbb{C}) \oplus \mathfrak{sl}_3(\mathbb{C})$ into an exceptional Lie algebra of type E_6, and deduced that the algebra of invariants for the natural representation of $\mathfrak{sl}_{333}(\mathbb{C})$ is freely generated by homogeneous polynomials in degrees 6, 9 and 12. A general computational framework for studying θ-groups has been developed recently by de Graaf [5].

An expression for the fundamental invariant of degree 9 in terms of 3×3 slices was obtained by Strassen [15, Lemma 4.5]. A geometric interpretation of Strassen's result has been given by Ottaviani [14, §3]; see also Domokos and Drensky [6].

Date: 28 March 2012.

2010 *Mathematics Subject Classification.* Primary 13A50; Secondary 15A72, 17B10.

Key words and phrases. Multidimensional arrays, polynomial invariants, semisimple Lie algebras, representation theory.

Nurmiev [13] obtained an implicit description of all three invariants in terms of convolutions of volume forms. An approach to these polynomials using classical invariant theory has been given by Briand et al. [4]. For an application to quantum information theory, see Duff and Ferrara [7]. For recent related work on $2\times 2\times 2\times 2$ arrays, see Huggins et al. [9].

In this paper we obtain explicit expressions for the fundamental invariants in degrees 6, 9, 12 for the action of $SL_3(\mathbb{C}) \times SL_3(\mathbb{C}) \times SL_3(\mathbb{C})$ on $3 \times 3 \times 3$ arrays. These invariants have respectively 1152, 9216 and 209061 terms. We express these homogeneous polynomials in terms of the orbits of the finite symmetry group $(S_3 \times S_3 \times S_3) \rtimes S_3$ acting on monomials of weight zero for the action of $\mathfrak{sl}_{333}(\mathbb{C})$. We use this action to efficiently represent polynomials as orbit sums; the invariants, especially in degree 12, involve far too many monomials to be written down completely. Using this method, it becomes possible to represent the fundamental invariants in a paper (and not just on a hard disk). Files containing the invariants are available as ancillary files attached to the `arXiv` version of this paper.

2. Preliminaries

2.1. **Homogeneous polynomials on $3\times 3\times 3$ arrays.** Let e_1, e_2, e_3 be the standard basis vectors of $\mathbb{C}^3$. We consider the tensor product $\mathbb{C}^{333} = \mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$. Every element of $\mathbb{C}^{333}$ is a finite sum of elements of the form $u \otimes v \otimes w$ where $u, v, w \in \mathbb{C}^3$. A basis for $\mathbb{C}^{333}$ consists of the 27 simple tensors $e_{ijk} = e_i \otimes e_j \otimes e_k$ for $1 \le i, j, k \le 3$. We therefore identify the array $X = (x_{ijk})$ with the element

$$X = \sum_{i=1}^{3}\sum_{j=1}^{3}\sum_{k=1}^{3} x_{ijk} e_{ijk}, \qquad x_{ijk} \in \mathbb{C}.$$

We introduce variables x_{ijk} for $1 \le i, j, k \le 3$ corresponding to the entries of X; this notation is ambiguous but should not be confusing. Strictly speaking, x_{ijk} is a coordinate function on $\mathbb{C}^{333}$ and is therefore a dual basis vector $e_i^* \otimes e_j^* \otimes e_k^*$, but this distinction will not be important for us. We consider the polynomial algebra $P = \mathbb{C}[\, x_{ijk} \mid 1 \le i, j, k \le 3 \,]$. A basis of P consists of the monomials

$$M(E) = \prod_{i=1}^{3}\prod_{j=1}^{3}\prod_{k=1}^{3} x_{ijk}^{e_{ijk}} = x_{111}^{e_{111}} \cdots x_{ijk}^{e_{ijk}} \cdots x_{333}^{e_{333}}, \tag{1}$$

where $E = (e_{ijk})$ is an exponent array of non-negative integers. The homogeneous subspace P_N of degree N has a basis consisting of monomials for which

$$\sum_{i=1}^{3}\sum_{j=1}^{3}\sum_{k=1}^{3} e_{ijk} = N.$$

A linear operator on $\mathbb{C}^3$ is represented by a 3×3 matrix $A = (a_{ij})$ where $a_{ij} \in \mathbb{C}$. The action of a triple of invertible operators (A, B, C) on x_{ijk} is given by

$$(A, B, C) \cdot x_{ijk} = \sum_{p=1}^{3}\sum_{q=1}^{3}\sum_{r=1}^{3} a_{pi} b_{qj} c_{rk} x_{pqr}.$$

This action extends to polynomials as follows:

$$(A, B, C) \cdot f\big(\dots, x_{ijk}, \dots\big) = f\big(\dots, (A, B, C) \cdot x_{ijk}, \dots\big).$$

We call f an invariant if $(A, B, C) \cdot f = f$ when $\det(A) = \det(B) = \det(C) = 1$.

2.2. **Representations of Lie algebras.** The $n \times n$ complex matrices of determinant 1 form the special linear group $SL_n(\mathbb{C})$. Finite-dimensional representations of $SL_n(\mathbb{C})$ can be studied in terms of the Lie algebra $\mathfrak{sl}_n(\mathbb{C})$, which consists of all $n \times n$ complex matrices of trace 0. The standard basis of $\mathfrak{sl}_n(\mathbb{C})$ consists of

- the matrix units $U_{i,j}$ for $i \neq j$ with (i,j) entry 1 and other entries 0,
- the diagonal matrices $H_i = U_{i,i} - U_{i+1,i+1}$ for $i = 1, 2, \dots, n{-}1$.

The simple root vectors are the matrix units $T_i = U_{i,i+1}$ for $i = 1, 2, \dots, n{-}1$. The natural representation of $\mathfrak{sl}_n(\mathbb{C})$ is its irreducible action on $\mathbb{C}^n$. We have

$$H_i \cdot e_j = \begin{cases} e_j & \text{if } j = i \\ -e_j & \text{if } j = i{+}1 \\ 0 & \text{otherwise,} \end{cases} \qquad T_i \cdot e_j = \begin{cases} e_{j-1} & \text{if } j = i{+}1 \\ 0 & \text{otherwise.} \end{cases}$$

We are concerned exclusively with the Lie algebra $\mathfrak{sl}_3(\mathbb{C})$; in this case

$$H_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \; H_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \; T_1 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \; T_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}.$$

The brackets of H_i and T_j are

$$(2) \quad [H_1, T_1] = 2T_1, \qquad [H_1, T_2] = -T_2, \qquad [H_2, T_1] = -T_1, \qquad [H_2, T_2] = 2T_2.$$

We consider the action of $\mathfrak{sl}_{333}(\mathbb{C}) = \mathfrak{sl}_3(\mathbb{C}) \oplus \mathfrak{sl}_3(\mathbb{C}) \oplus \mathfrak{sl}_3(\mathbb{C})$ on its irreducible representation $\mathbb{C}^{333}$. For $\ell = 1, 2, 3$ we write a superscript (ℓ) to indicate the basis elements of the ℓ-th summand. These elements act on x_{ijk} as follows $(m = 1, 2)$:

$$H^{(1)}_m \cdot x_{ijk} = \begin{cases} x_{ijk} & \text{if } i = m \\ -x_{ijk} & \text{if } i = m{+}1 \\ 0 & \text{otherwise} \end{cases} \qquad T^{(1)}_m \cdot x_{ijk} = \begin{cases} x_{i-1,j,k} & \text{if } i = m{+}1 \\ 0 & \text{otherwise} \end{cases}$$

$$H^{(2)}_m \cdot x_{ijk} = \begin{cases} x_{ijk} & \text{if } j = m \\ -x_{ijk} & \text{if } j = m{+}1 \\ 0 & \text{otherwise} \end{cases} \qquad T^{(2)}_m \cdot x_{ijk} = \begin{cases} x_{i,j-1,k} & \text{if } j = m{+}1 \\ 0 & \text{otherwise} \end{cases}$$

$$H^{(3)}_m \cdot x_{ijk} = \begin{cases} x_{ijk} & \text{if } k = m \\ -x_{ijk} & \text{if } k = m{+}1 \\ 0 & \text{otherwise} \end{cases} \qquad T^{(3)}_m \cdot x_{ijk} = \begin{cases} x_{i,j,k-1} & \text{if } k = m{+}1 \\ 0 & \text{otherwise} \end{cases}$$

The action of a Lie algebra L on a tensor product $V \otimes W$ of L-modules is given by $A \cdot (v \otimes w) = (A \cdot v) \otimes w + v \otimes (A \cdot w)$ for $A \in L$, $v \in V$, $w \in W$. If we identify the N-th symmetric power $S^N V$ of the p-dimensional L-module V with the space of homogeneous polynomials of degree N on a basis $v_1, \dots, v_p$ of V, then the action of L on $S^N V$ is given by

$$\begin{aligned} A \cdot (v_1^{e_1} v_2^{e_2} \cdots v_p^{e_p}) &= \sum_{i=1}^{p} v_1^{e_1} \cdots (A \cdot v_i^{e_i}) \cdots v_p^{e_p} \\ &= \sum_{i=1}^{p} v_1^{e_1} \cdots \big(e_i v_i^{e_i - 1} (A \cdot v_i) \big) \cdots v_p^{e_p} = \sum_{i=1}^{p} e_i \, v_1^{e_1} \cdots v_i^{e_i - 1} \cdots v_p^{e_p} (A \cdot v_i). \end{aligned}$$

Lemma 2.1. *For $m = 1, 2$ we have*

$$H_m^{(1)} \cdot \big(x_{111}^{e_{111}} \cdots x_{333}^{e_{333}} \big) = \sum_{j=1}^{3} \sum_{k=1}^{3} (e_{m,j,k} - e_{m+1,j,k}) \, x_{111}^{e_{111}} \cdots x_{333}^{e_{333}},$$

$$H_m^{(2)} \cdot \big(x_{111}^{e_{111}} \cdots x_{333}^{e_{333}} \big) = \sum_{i=1}^{3} \sum_{k=1}^{3} (e_{i,m,k} - e_{i,m+1,k}) \, x_{111}^{e_{111}} \cdots x_{333}^{e_{333}},$$

$$H_m^{(3)} \cdot \big(x_{111}^{e_{111}} \cdots x_{333}^{e_{333}} \big) = \sum_{i=1}^{3} \sum_{j=1}^{3} (e_{i,j,m} - e_{i,j,m+1}) \, x_{111}^{e_{111}} \cdots x_{333}^{e_{333}}.$$

The eigenvalues of $x_{111}^{e_{111}} \cdots x_{333}^{e_{333}}$ for the elements $H_m^{(\ell)}$ will be denoted

$$\omega_{11} = \sum_{j=1}^{3} \sum_{k=1}^{3} (e_{1jk} - e_{2jk}), \quad \omega_{12} = \sum_{j=1}^{3} \sum_{k=1}^{3} (e_{2jk} - e_{3jk}), \quad \omega_{21} = \sum_{i=1}^{3} \sum_{k=1}^{3} (e_{i1k} - e_{i2k}),$$

$$\omega_{22} = \sum_{i=1}^{3} \sum_{k=1}^{3} (e_{i2k} - e_{i3k}), \quad \omega_{31} = \sum_{i=1}^{3} \sum_{j=1}^{3} (e_{ij1} - e_{ij2}), \quad \omega_{32} = \sum_{i=1}^{3} \sum_{j=1}^{3} (e_{ij2} - e_{ij3}).$$

The weight of $x_{111}^{e_{111}} \cdots x_{333}^{e_{333}}$ is the tuple $(\omega_{11}, \omega_{12}, \omega_{21}, \omega_{22}, \omega_{31}, \omega_{32})$. A monomial has weight zero if $\omega_{\ell m} = 0$ for $\ell = 1, 2, 3$ and $m = 1, 2$. For given degree and given weight, the weight space $W(N \mid \omega_{11}, \omega_{12}, \omega_{21}, \omega_{22}, \omega_{31}, \omega_{32})$ is the subspace of P_N spanned by the corresponding monomials.

Lemma 2.2. *For $m = 1, 2$ we have*

$$T_m^{(1)} \cdot \big(x_{111}^{e_{111}} \cdots x_{333}^{e_{333}} \big) = \sum_{j=1}^{3} \sum_{k=1}^{3} e_{m+1,j,k} \, x_{111}^{e_{111}} \cdots x_{m,j,k}^{e_{m,j,k}+1} \cdots x_{m+1,j,k}^{e_{m+1,j,k}-1} \cdots x_{333}^{e_{333}},$$

$$T_m^{(2)} \cdot \big(x_{111}^{e_{111}} \cdots x_{333}^{e_{333}} \big) = \sum_{i=1}^{3} \sum_{k=1}^{3} e_{i,m+1,k} \, x_{111}^{e_{111}} \cdots x_{i,m,k}^{e_{i,m,k}+1} \cdots x_{i,m+1,k}^{e_{i,m+1,k}-1} \cdots x_{333}^{e_{333}},$$

$$T_m^{(3)} \cdot \big(x_{111}^{e_{111}} \cdots x_{333}^{e_{333}} \big) = \sum_{i=1}^{3} \sum_{j=1}^{3} e_{i,j,m+1} \, x_{111}^{e_{111}} \cdots x_{i,j,m}^{e_{i,j,m}+1} \cdots x_{i,j,m+1}^{e_{i,j,m+1}-1} \cdots x_{333}^{e_{333}}.$$

By (2) the elements $T_m^{(\ell)}$ induce linear maps:

$$\begin{aligned}
T_1^{(1)} &\colon W(N \mid 0,0,0,0,0,0) \longrightarrow W(N \mid 2,-1,0,0,0,0), \\
T_2^{(1)} &\colon W(N \mid 0,0,0,0,0,0) \longrightarrow W(N \mid -1,2,0,0,0,0), \\
T_1^{(2)} &\colon W(N \mid 0,0,0,0,0,0) \longrightarrow W(N \mid 0,0,2,-1,0,0), \\
T_2^{(2)} &\colon W(N \mid 0,0,0,0,0,0) \longrightarrow W(N \mid 0,0,-1,2,0,0), \\
T_1^{(3)} &\colon W(N \mid 0,0,0,0,0,0) \longrightarrow W(N \mid 0,0,0,0,2,-1), \\
T_2^{(3)} &\colon W(N \mid 0,0,0,0,0,0) \longrightarrow W(N \mid 0,0,0,0,-1,2).
\end{aligned}$$

We write $\Omega_{\ell m}$ for the nonzero weights and form the direct sum:

$$\Lambda_N = (T_1^{(1)}, \dots, T_2^{(3)}) \colon W(N \mid 0,0,0,0,0,0) \longrightarrow \bigoplus_{\ell=1}^{3} \bigoplus_{m=1}^{2} W(N \mid \Omega_{\ell m}).$$

Lemma 2.3. *The invariant polynomials in degree N for the action of $\mathfrak{sl}_{333}(\mathbb{C})$ on the space $\mathbb{C}^{333}$ are the (nonzero) elements of the kernel of the linear map Λ_N.*

Theorem 2.4. *The algebra of invariants for* $\mathfrak{sl}_{333}(\mathbb{C})$ *acting on* $\mathbb{C}^{333}$ *is freely generated by three fundamental invariants in degrees* 6, 9 *and* 12*. Thus for* $N = 6$, 9, 12 *the nullspace of* Λ_N *has dimension* 1, 1, 2 *respectively.*

Proof. Vinberg [16]; see especially item 2 in the table on page 491. □

2.3. **Combinatorics of monomials.** Let $E = (e_{ijk})$ be a $3 \times 3 \times 3$ exponent array with non-negative integer entries corresponding to the monomial (1).

Definition 2.5. The **flattening** of E is the ordered list obtained by applying lexicographical order to the subscripts:

$$\text{flatten}(E) = [e_{111}, e_{112}, e_{113}, e_{121}, e_{122}, e_{123}, \dots, e_{331}, e_{332}, e_{333}].$$

The **total order** on exponent arrays is the lex order on flattenings: that is, $E < E'$ for $\text{flatten}(E) = [f_1, f_2, \dots, f_{27}]$ and $\text{flatten}(E') = [f'_1, f'_2, \dots, f'_{27}]$ if and only if $f_i < f'_i$ where i is the least index with $f_i \neq f'_i$. The **matrix form** of E is:

$$\left[\begin{array}{ccc|ccc|ccc} e_{111} & e_{121} & e_{131} & e_{112} & e_{122} & e_{132} & e_{113} & e_{123} & e_{133} \\ e_{211} & e_{221} & e_{231} & e_{212} & e_{222} & e_{232} & e_{213} & e_{223} & e_{233} \\ e_{311} & e_{321} & e_{331} & e_{212} & e_{222} & e_{232} & e_{213} & e_{223} & e_{233} \end{array} \right].$$

The third index distinguishes the frontal slices, where a **slice** is a 3×3 matrix obtained by fixing one subscript; there are three parallel slices in each direction. We call E an **equal parallel slice array** if parallel slices have the same sum.

Lemma 2.6. *A basis for* $W(N \mid 0,0,0,0,0,0)$ *consists of the monomials of degree* N *whose exponent arrays are equal parallel slice arrays. In degree* N*, there are no monomials of weight zero, and hence no invariants, unless* N *is a multiple of* 3*.*

Proof. This follows immediately from Lemma 2.1. □

To generate the exponent arrays for weight zero monomials of degree N, we use nested loops and the condition that the horizontal parallel slices have sum $N/3$. (See Table 1.) To generate the higher weight monomials, we first use Lemma 2.2 to generate the monomials for weights Ω_{1m} ($m = 1, 2$). We then use symmetry to obtain the monomials for weights $\Omega_{\ell m}$ ($\ell = 2, 3$, $m = 1, 2$): if E is an exponent array of weight Ω_{1m} then E' defined by $e'_{ijk} = e_{jik}$ is an exponent array of weight Ω_{2m}, and similarly for monomials of weight Ω_{3m}. (See Table 2.)

Definition 2.7. The **symmetry group** acting on the weight zero monomials in degree N is the semidirect product $G = (S_3 \times S_3 \times S_3) \rtimes S_3$. Each factor in $S_3 \times S_3 \times S_3$ permutes the parallel slices in the corresponding direction; the last S_3 permutes the directions. More precisely, (α, β, γ) and δ act on $E = (e_{ijk})$ by:

$$\big((\alpha, \beta, \gamma) \cdot E\big)_{i,j,k} = e_{\alpha(i), \beta(j), \gamma(k)}, \qquad \big(\delta \cdot E\big)_{i,j,k} = e_{i^\delta, j^\delta, k^\delta}.$$

The **orbit** of a weight zero monomial $M(E)$ is $\mathcal{O}(M) = \{\, M(g \cdot E) \mid g \in G \,\}$. The **symmetric** and **alternating orbit sums** are defined by

$$\mathcal{O}^+(M) = \frac{|\mathcal{O}(M)|}{|G|} \sum_{g \in G} M(g \cdot E), \qquad \mathcal{O}^-(M) = \frac{|\mathcal{O}(M)|}{|G|} \sum_{g \in G} \epsilon(g)\, M(g \cdot E),$$

where $\epsilon(g)$ is the product of the signs of the components of $g = (\alpha, \beta, \gamma, \delta) \in G$. For some monomials, the alternating orbit sum will be zero.

See Table 3 for procedures to generate the orbit of a weight zero monomial; `smallorbit` permutes the slices, and `largeorbit` permutes the directions.

```
• set weightzeromonomials ← [ ] (empty list)
• for $f_1$ from 0 to $N/3$ do for $f_2$ from 0 to $N/3 - f_1$ do ...
  for $f_8$ from 0 to $N/3 - (f_1 + \cdots + f_7)$ do:
    · set $f_9 \leftarrow N/3 - (f_1 + \cdots + f_8)$
    · for $f_{10}$ from 0 to $N/3$ do for $f_{11}$ from 0 to $N/3 - f_{10}$ do ...
      for $f_{17}$ from 0 to $N/3 - (f_{10} + \cdots + f_{16})$ do:
        · set $f_{18} \leftarrow N/3 - (f_{10} + \cdots + f_{17})$
        · for $f_{19}$ from 0 to $N/3$ do for $f_{20}$ from 0 to $N/3 - f_{19}$ do ...
          for $f_{26}$ from 0 to $N/3 - (f_{19} + \cdots + f_{25})$ do:
            · set $f_{27} \leftarrow N/3 - (f_{19} + \cdots + f_{26})$
            · set $e \leftarrow$ unflatten($f$)
            · if $\sum_{j,k} e_{1jk} = \sum_{j,k} e_{2jk} = \sum_{j,k} e_{3jk}$ and
              $\sum_{i,k} e_{i1k} = \sum_{i,k} e_{i2k} = \sum_{i,k} e_{i3k}$ and
              $\sum_{i,j} e_{ij1} = \sum_{i,j} e_{ij2} = \sum_{i,j} e_{ij3}$ then
              append $[f_1, \dots, f_{27}]$ to weightzeromonomials
• return weightzeromonomials
```

Table 1. Pseudocode to generate weight zero monomials of degree N

```
• set $\ell \leftarrow 1$
• for $m = 1, 2$ do
    · set higherweightmonomials[$\ell, m$] ← [ ] (empty list)
    · for $x$ in weightzeromonomials do
        · set $e \leftarrow$ unflatten($x$)
        · for $j = 1, 2, 3$ do for $k = 1, 2, 3$ do
            · if $e_{m+1,j,k} > 0$ then
                set $f \leftarrow e$
                set $f_{m+1,j,k} \leftarrow f_{m+1,j,k} - 1$
                set $f_{m,j,k} \leftarrow f_{m,j,k} + 1$
                append flatten($f$) to higherweightmonomials[$\ell, m$]
• return higherweightmonomials[$\ell, 1$], higherweightmonomials[$\ell, 2$]
```

Table 2. Pseudocode to generate higher weight monomials ($\ell = 1$)

3. Degree 6

Lemma 3.1. *In degree* 6*, there are* 1152 *monomials of weight zero, and* 792 *monomials of each higher weight* $\Omega_{\ell m}$ *for* $\ell = 1, 2, 3$ *and* $m = 1, 2$.

Proof. We generate the monomials using an implementation in Maple 15 of the algorithms in Tables 1 and 2 with $N = 6$. □

Lemma 3.2. *The nullspace of the matrix representing* Λ_6 *has dimension* 1 *modulo* $p = 101$. *Using symmetric representatives, the canonical basis vector of the nullspace has coefficients* $\{-10, -4, -2, 1, 2, 4, 8\}$. *If we interpret these as integers then the corresponding polynomial is an invariant for the action of* $\mathfrak{sl}_{333}(\mathbb{C})$.

Proof. We use the `LinearAlgebra[Modular]` package in Maple 15 to create a matrix B with an upper block of size 1152×1152 and a lower block of size 792×1152.

```
• smallgrouporbit($f$)
    · set $e \leftarrow$ unflatten($f$)
    · set orbit $\leftarrow \{\,\}$
    · for $p \in S_3$ do for $q \in S_3$ do for $r \in S_3$ do:
        · for $i = 1,2,3$ do for $j = 1,2,3$ do for $k = 1,2,3$ do:
            set $f_{i,j,k} \leftarrow e_{p(i),q(j),r(k)}$
        · set orbit $\leftarrow$ orbit $\cup$ {flatten($f$)}
    · return orbit

• largegrouporbit($f$)
    · set $e \leftarrow$ unflatten($f$)
    · set orbit $\leftarrow \{\,\}$
    · for $s \in S_3$ do:
        · for $i = 1,2,3$ do for $j = 1,2,3$ do for $k = 1,2,3$ do:
            set $t \leftarrow (i, j, k)$
            set $f_{i,j,k} \leftarrow e_{t_{s(1)},t_{s(2)},t_{s(3)}}$
        · orbit $\leftarrow$ orbit $\cup$ smallgrouporbit(flatten($f$))
    · return orbit
```

Table 3. Pseudocode to generate the orbits of a weight zero monomial

```
• set $B \leftarrow$ Matrix$(1944, 1152)$ modulo 101
• set $\ell \leftarrow 1$
• set $c \leftarrow 0$
• for $x \in$ weightzeromonomials do
    · set $c \leftarrow c + 1$
    · for $m = 1, 2$ do
        · set $e \leftarrow$ unflatten($x$)
        · for $j = 1,2,3$ do for $k = 1,2,3$ do
            · if $e_{m+1,j,k} > 0$ then
                set $f \leftarrow e$
                set $f_{m+1,j,k} \leftarrow f_{m+1,j,k} - 1$
                set $f_{m,j,k} \leftarrow f_{m,j,k} + 1$
                set $g \leftarrow$ flatten($f$)
                set $r \leftarrow$ monomialindex($g, \ell, m$):
                set $B_{1152+r,c} \leftarrow (B_{1152+r,c} + e_{m+1,j,k})$ modulo 101
• compute the row canonical form of $B$
```

Table 4. Pseudocode to fill the matrix ($N = 6$, $\ell = 1$)

For $\ell = 1, 2, 3$ and $m = 1, 2$ we apply Lemma 2.2 to store the matrix representing $T_m^{(\ell)}$ in the lower block; we then compute the row canonical form. See Table 4 for $\ell = 1$; `monomialindex` does a binary search for a higher weight monomial in the lexicographically ordered list of all monomials of that weight. At termination, B has rank 1151. From the row canonical form, we extract a basis vector for the nullspace. We then perform another computation using integer arithmetic to verify that the corresponding polynomial is indeed an invariant over $\mathbb{C}$. □

<table>
<tr><th>I_6</th><th>MINIMAL REPRESENTATIVE</th><th>ORBIT SIZE</th><th>#</th></tr>
<tr><td>-10</td><td>$\left[\begin{array}{ccc|ccc|ccc} 0&0&0&0&0&1&0&1&0\\ 0&0&1&0&0&0&1&0&0\\ 0&1&0&1&0&0&0&0&0 \end{array}\right]$</td><td>36</td><td>1</td></tr>
<tr><td>-4</td><td>$\left[\begin{array}{ccc|ccc|ccc} 0&0&0&0&0&1&0&0&1\\ 0&1&0&0&0&0&1&0&0\\ 1&0&0&0&1&0&0&0&0 \end{array}\right]$</td><td>324</td><td>2</td></tr>
<tr><td>-2</td><td>$\left[\begin{array}{ccc|ccc|ccc} 0&0&0&0&0&0&0&0&2\\ 0&1&0&0&1&0&0&0&0\\ 1&0&0&1&0&0&0&0&0 \end{array}\right]$</td><td>162</td><td>3</td></tr>
<tr><td>1</td><td>$\left[\begin{array}{ccc|ccc|ccc} 0&0&0&0&0&0&0&0&2\\ 0&0&0&0&2&0&0&0&0\\ 2&0&0&0&0&0&0&0&0 \end{array}\right]$</td><td>36</td><td>4</td></tr>
<tr><td>2</td><td>$\left[\begin{array}{ccc|ccc|ccc} 0&0&0&0&0&1&0&0&1\\ 0&1&0&0&0&0&0&1&0\\ 1&0&0&1&0&0&0&0&0 \end{array}\right]$</td><td>108</td><td>5</td></tr>
<tr><td>2</td><td>$\left[\begin{array}{ccc|ccc|ccc} 0&0&0&0&0&1&0&0&1\\ 0&0&0&0&1&0&1&0&0\\ 1&1&0&0&0&0&0&0&0 \end{array}\right]$</td><td>324</td><td>6</td></tr>
<tr><td>4</td><td>$\left[\begin{array}{ccc|ccc|ccc} 0&0&0&0&0&0&0&0&2\\ 0&1&0&1&0&0&0&0&0\\ 1&0&0&0&1&0&0&0&0 \end{array}\right]$</td><td>54</td><td>7</td></tr>
<tr><td>8</td><td>$\left[\begin{array}{ccc|ccc|ccc} 0&0&0&0&0&1&0&1&0\\ 0&0&1&0&0&0&1&0&0\\ 1&0&0&0&1&0&0&0&0 \end{array}\right]$</td><td>108</td><td>8</td></tr>
</table>

TABLE 5. The fundamental invariant in degree 6

Lemma 3.3. *The coefficients of the canonical basis vector for the nullspace of* Λ_6 *are constant on orbits for the action of* $(S_3 \times S_3 \times S_3) \rtimes S_3$ *on weight zero monomials. For each orbit, the coefficient, the matrix form of the minimal representative, and the orbit size, are displayed in Table* 5.

Proof. We implemented the algorithms of Tables 3 and 4 in Maple 15. □

Theorem 3.4. *Every invariant in degree* 6 *for* $SL_3(\mathbb{C}) \times SL_3(\mathbb{C}) \times SL_3(\mathbb{C})$ *acting on* $3 \times 3 \times 3$ *arrays* $X = (x_{ijk})$ *is a scalar multiple of*

$$
\begin{aligned}
&- 10\, \mathcal{O}^+\big(x_{123}x_{132}x_{213}x_{231}x_{312}x_{321} \big) - 4\, \mathcal{O}^+\big(x_{132}x_{133}x_{213}x_{221}x_{311}x_{322} \big) \\
&- 2\, \mathcal{O}^+\big(x_{133}^2 x_{221}x_{222}x_{311}x_{312} \big) + \mathcal{O}^+\big(x_{133}^2 x_{222}^2 x_{311}^2 \big) \\
&+ 2\, \mathcal{O}^+\big(x_{132}x_{133}x_{221}x_{223}x_{311}x_{312} \big) + 2\, \mathcal{O}^+\big(x_{132}x_{133}x_{213}x_{222}x_{311}x_{321} \big) \\
&+ 4\, \mathcal{O}^+\big(x_{133}^2 x_{212}x_{221}x_{311}x_{322} \big) + 8\, \mathcal{O}^+\big(x_{123}x_{132}x_{213}x_{231}x_{311}x_{322} \big).
\end{aligned}
$$

Proof. This is a restatement of the results in Table 5. □

4. DEGREE 9

Lemma 4.1. *In degree* 9*, there are* 22620 *monomials of weight zero, and* 17802 *monomials of each higher weight* $\Omega_{\ell m}$ *for* $\ell = 1, 2, 3$ *and* $m = 1, 2$.

Proof. Similar to the proof of Lemma 3.1. □

Lemma 4.2. *The nullspace of the matrix representing* Λ_9 *has dimension* 1 *modulo* $p = 101$. *Using symmetric representatives, the canonical basis vector of the nullspace has coefficients* $\{-1, 0, 1\}$*; the nonzero coefficients each occur* 4608 *times*

I_9	MINIMAL REPRESENTATIVE	ORBIT SIZE	#
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 0&1&0&0&0&1&1&0&0\\ 2&0&0&0&1&0&0&0&0 \end{array}\right]$	648	1
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 0&1&1&0&0&0&1&0&0\\ 0&1&0&2&0&0&0&0&0 \end{array}\right]$	648	2
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 0&1&0&0&1&0&1&0&0\\ 1&0&1&1&0&0&0&0&0 \end{array}\right]$	1296	3
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 0&2&0&0&0&0&1&0&0\\ 1&0&0&1&0&1&0&0&0 \end{array}\right]$	648	4
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 0&0&1&1&0&0&0&1&0\\ 1&1&0&1&0&0&0&0&0 \end{array}\right]$	648	5
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 0&0&1&1&1&0&0&0&0\\ 2&0&0&0&0&0&0&1&0 \end{array}\right]$	1296	6
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 0&1&0&1&0&1&0&0&0\\ 1&1&0&0&0&0&1&0&0 \end{array}\right]$	648	7
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 0&1&1&1&0&0&0&0&0\\ 1&0&0&0&1&0&1&0&0 \end{array}\right]$	1296	8
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 0&1&0&1&0&0&1&0&0\\ 0&1&1&1&0&0&0&0&0 \end{array}\right]$	216	9
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 0&0&1&2&0&0&0&0&0\\ 0&2&0&0&0&0&1&0&0 \end{array}\right]$	648	10
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&1&0&1&1\\ 1&0&0&1&0&1&0&0&0\\ 1&1&0&0&0&0&0&1&0 \end{array}\right]$	72	11
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&2&0&1&0\\ 0&1&1&0&0&0&1&0&0\\ 1&0&0&1&0&0&0&1&0 \end{array}\right]$	432	12
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&2&0&1&0\\ 0&1&0&0&1&0&1&0&0\\ 2&0&0&0&0&0&0&0&1 \end{array}\right]$	648	13
1	$\left[\begin{array}{ccc\|ccc\|ccc} 0&0&0&0&0&2&0&1&0\\ 0&0&1&0&0&0&2&0&0\\ 0&2&0&1&0&0&0&0&0 \end{array}\right]$	72	14

TABLE 6. The fundamental invariant in degree 9

and 0 *occurs* 13404 *times. If we interpret these as integers then the corresponding polynomial is an invariant for the action of* $\mathfrak{sl}_{333}(\mathbb{C})$.

Proof. Similar to the proof of Lemma 3.2. ☐

Lemma 4.3. *The canonical basis vector for the nullspace of* Λ_9 *is a linear combination of alternating orbit sums for the action of* $(S_3 \times S_3 \times S_3) \rtimes S_3$ *on weight zero monomials. For each orbit, the coefficient, the matrix form of the minimal representative, and the orbit size, are displayed in Table* 6.

Proof. Similar to the proof of Lemma 3.3. (There are another 30 orbits for the action of the symmetry group but they all occur with coefficient 0.) ☐

Theorem 4.4. *Every invariant in degree* 9 *for* $SL_3(\mathbb{C}) \times SL_3(\mathbb{C}) \times SL_3(\mathbb{C})$ *acting on* $3 \times 3 \times 3$ *arrays* $X = (x_{ijk})$ *is a scalar multiple of*

$$
\begin{aligned}
&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{213}x_{221}x_{232}x_{311}^2x_{322}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{213}x_{221}x_{231}x_{312}^2x_{321}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{213}x_{221}x_{222}x_{311}x_{312}x_{331}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{213}x_{221}^2x_{311}x_{312}x_{332}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{212}x_{223}x_{231}x_{311}x_{312}x_{321}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{212}x_{222}x_{231}x_{311}^2x_{323}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{212}x_{221}x_{232}x_{311}x_{313}x_{321}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{212}x_{221}x_{231}x_{311}x_{313}x_{322}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{212}x_{213}x_{221}x_{312}x_{321}x_{331}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{212}^2x_{231}x_{313}x_{321}^2\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}x_{133}x_{211}x_{212}x_{232}x_{311}x_{321}x_{323}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}^2x_{213}x_{221}x_{231}x_{311}x_{312}x_{323}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}^2x_{213}x_{221}x_{222}x_{311}^2x_{333}\big) \\
+&\mathcal{O}^-\big(x_{123}x_{132}^2x_{213}^2x_{231}x_{312}x_{321}^2\big).
\end{aligned}
$$

Proof. This is a restatement of the results in Table 6. □

5. DEGREE 12

We find explicit forms of two linearly independent invariants in degree 12, and verify that the simpler invariant can be taken as the generator in degree 12.

Lemma 5.1. *In degree* 12*, there are* 302274 *monomials of weight zero, and* 254961 *monomials of each higher weight* $\Omega_{\ell m}$ *for* $\ell = 1, 2, 3$ *and* $m = 1, 2$*.*

Even using modular arithmetic, it is impossible to process efficiently a matrix with 302274 columns and $302274 + 254961 = 557235$ rows. Therefore we look for invariants that are linear combinations of the symmetric orbit sums.

Lemma 5.2. *In degree* 12*, there are* 359 *orbits for the action of* $(S_3 \times S_3 \times S_3) \rtimes S_3$ *on weight zero monomials. For each orbit, the flattened minimal representative and the orbit size are given in Tables* 7*–*11*.*

We consider a matrix with 359 columns, one for each symmetric orbit sum.

Lemma 5.3. *The nullspace of the matrix representing the restriction of* Λ_{12} *to the span of the symmetric orbit sums has dimension* 2 *modulo* $p = 101$*.*

Proof. We use modular arithmetic on a matrix M with an upper block of size 359×359 and a lower block of size 254961×359. For each $\ell = 1, 2, 3$ and $m = 1, 2$ we consider the restriction of $T_m^{(\ell)}$ to the subspace of $W(12 \mid 0, 0, 0, 0, 0, 0)$ spanned by the symmetric orbit sums; we store the corresponding matrix in the lower block, and compute the row canonical form. After the first iteration ($\ell = m = 1$) the rank is 357, and does not increase for the remaining five iterations. □

Lemma 5.3 agrees with Theorem 2.4: a basis of the invariants in degree 12 consists of I_6^2 together with a new generator I_{12}. We confirm this result with further computations using integer arithmetic.

Lemma 5.4. *Let A be a matrix with integer entries. If r_0 is its rank over $\mathbb{Q}$ and r_p its its rank over the prime field $\mathbb{F}_p$, then $r_p \le r_0$. Hence the dimension of the nullspace over $\mathbb{Q}$ is no larger than the dimension of the nullspace over $\mathbb{F}_p$.*

Proof. The rank of a matrix is r if and only if all $(r+1)$-minors are zero, and at least one r-minor is not zero. Since these minors are polynomials in the matrix entries with integer coefficients, the claim follows. $\square$

Theorem 5.5. *In degree 12, the polynomials I_{12} and I'_{12} given in Tables 7–11 form a reduced basis for the lattice of invariants with integer coefficients.*

Proof. For each $\ell = 1, 2, 3$ and $m = 1, 2$ we totally order the 254961 higher weight monomials, and partition each set into 399 blocks of 639 consecutive monomials. For $j = 1, 2, \ldots 359$ we compute the action of $T_m^{(\ell)}$ on the j-th symmetric orbit sum, and separate the resulting terms corresponding to the different blocks.

We create an integer matrix M with an upper block of size 359×359 and a lower block of size 639×359, initialized to zero. For $\ell = 1, 2, 3$, $m = 1, 2$, $j = 1, 2, \ldots, 359$, $k = 1, 2, \ldots, 399$ we consider the terms in block k obtained by applying $T_m^{(\ell)}$ to the j-th symmetric orbit sum. We store the integer coefficients of these higher weight monomials in the lower block, and compute the Hermite normal form (HNF).

We obtain the same behavior as in Lemma 5.3: after the first iteration the rank is 357, and does not increase for the remaining iterations. At termination, we have an integer matrix of size 357×359, also called M, whose integer nullspace consists of the coefficient vectors of the invariant polynomials of degree 12.

To find a lattice basis for this integer nullspace, we compute the HNF of the transpose M^t, obtaining integer matrices U (invertible) and H, of sizes 359×359 and 359×357 respectively, for which $UM^t = H$. The last two rows of U form a lattice basis for the integer nullspace of M. We apply the algorithm of Gauss-Lagrange to these last two rows to find a short basis of the integer nullspace. (For an introduction to lattice basis reduction, see Bremner [1].) Since the lattice is 2-dimensional, the output consists of a shortest nonzero vector and a shortest vector which is not a multiple of the first vector. The components of these reduced basis vectors I_{12} and I'_{12} are given in Tables 7–11. Finally, we perform an independent check using integer arithmetic that I_{12} and I'_{12} are annihilated by all the $T_m^{(\ell)}$. $\square$

Lemma 5.6. *We have $I'_{12} = I_6^2 + 21\, I_{12}$. Hence I_{12} is the fundamental invariant in degree 12; it involves 235 of the 359 orbits, with a total of 209061 monomials.*

Proof. We compute the expression for I_6^2 as a linear combination of I_{12} and I'_{12}. This can be performed very efficiently in Maple 15 using the algorithms of Monagan and Pearce [11, 12] which are based on the work of Johnson [10]. $\square$

We can now state our main result, and an open problem.

Theorem 5.7. *The three fundamental invariants for $3 \times 3 \times 3$ arrays are:*

- *the polynomial I_6 of Table 5, which is a linear combination of the 8 symmetric orbits in degree 6, and has altogether 1152 terms;*

- *the polynomial* I_9 *of Table* 6*, which is the sum of* 14 *alternating orbits in degree* 9*, and has altogether* 9216 *terms;*
- *the polynomial* I_{12} *of Tables* 7–11*, which is a linear combination of* 235 *symmetric orbits in degree* 12*, and has altogether* 209061 *terms.*

Open Problem. Corollary 3.9 of Gelfand et al. [8] implies that the hyperdeterminant of a $3 \times 3 \times 3$ array has degree 36, and hence has the form

$$a\, I_6^6 + b\, I_6^4 I_{12} + c\, I_6^3 I_9^2 + d\, I_6^2 I_{12}^2 + e\, I_6 I_9^2 I_{12} + f\, I_9^4 + g\, I_{12}^3,$$

for some $a, b, c, d, e, f, g \in \mathbb{C}$. Determine these coefficients.

ACKNOWLEDGEMENTS

The first author was partially supported by a Discovery Grant from NSERC. The authors thank the referees and Luke Oeding for helpful comments.

REFERENCES

[1] M. R. Bremner: *Lattice Basis Reduction: An Introduction to the LLL Algorithm and its Applications*. CRC Press, 2012.
[2] M. R. Bremner: On the hyperdeterminant for $2 \times 2 \times 3$ arrays. *Linear and Multilinear Algebra* (available online 25 November 2011).
[3] M. R. Bremner, M. G. Bickis, M. Soltanifar: Cayley's hyperdeterminant: a combinatorial approach via representation theory. *Linear Algebra and Its Applications* (available online 1 March 2012).
[4] E. Briand, J.-G. Luque, J.-Y. Thibon, F. Verstraete: The moduli space of three-qutrit states. *Journal of Mathematical Physics* 45:12 (2004) 4855–4867.
[5] W. A. de Graaf: Computing representatives of nilpotent orbits of θ-groups. *Journal of Symbolic Computation* 46:4 (2011) 438–458.
[6] M. Domokos, V. Drensky: Defining relation for semi-invariants of three by three matrix triples. `arXiv:1101.3178v1 [math.RT]` (submitted 17 Jan 2011)
[7] M. J. Duff, S. Ferrara: E_6 and the bipartite entanglement of three qutrits. *Physical Review D* 76:12 (2007) 124023, 7 pages.
[8] I. M. Gelfand, M. M. Kapranov, A. V. Zelevinsky: Hyperdeterminants. *Advances in Mathematics* 96:2 (1992) 226–263.
[9] P. Huggins, B. Sturmfels, J. Yu, D. S. Yuster: The hyperdeterminant and triangulations of the 4-cube. *Mathematics of Computation* 77:263 (2008) 1653–1679.
[10] S. C. Johnson: Sparse polynomial arithmetic. *ACM SIGSAM Bulletin* 8:3 (1974) 63–71.
[11] M. Monagan, R. Pearce: Parallel sparse polynomial multiplication using heaps. *ISSAC 2009: Proceedings of the 2009 International Symposium on Symbolic and Algebraic Computation*, pages 263–269. ACM, 2009.
[12] M. Monagan, R. Pearce: Sparse polynomial division using a heap. *Journal of Symbolic Computation* 46:7 (2011) 807–822.
[13] A. G. Nurmiev: Orbits and invariants of third-order matrices. *Sbornik Mathematics* 191:5-6 (2000) 717–724.
[14] G. Ottaviani: Symplectic bundles on the plane, secant varieties and Lüroth quartics revisited. `arXiv:math/0702151v1 [math.AG]` (submitted 3 February 2007).
[15] V. Strassen: Rank and optimal computation of generic tensors. *Linear Algebra and Its Applications* 52/53 (1983) 345–685.
[16] E. B. Vinberg: The Weyl group of a graded Lie algebra. *Izvestiya Akademii Nauk SSSR, Seriya Matematicheskaya* 40:3 (1976) 488–526.

Department of Mathematics and Statistics, University of Saskatchewan, Canada
E-mail address: `bremner@math.usask.ca`

Department of Mathematics, Simon Fraser University, Canada
E-mail address: `hujiaxiong@gmail.com`

I_{12}	I'_{12}	MINIMAL REPRESENTATIVE	ORBIT SIZE	#
0	1	0 0 0 0 0 0 0 0 4 0 0 0 0 4 0 0 0 0 4 0 0 0 0 0 0 0 0	36	1
0	−4	0 0 0 0 0 0 0 0 4 0 0 0 1 3 0 0 0 0 3 1 0 0 0 0 0 0 0	324	2
0	6	0 0 0 0 0 0 0 0 4 0 0 0 2 2 0 0 0 0 2 2 0 0 0 0 0 0 0	162	3
0	4	0 0 0 0 0 0 0 0 4 0 1 0 1 2 0 0 0 0 2 1 0 1 0 0 0 0 0	324	4
0	8	0 0 0 0 0 0 0 0 4 0 1 0 1 2 0 0 0 0 3 0 0 0 1 0 0 0 0	216	5
0	−16	0 0 0 0 0 0 0 0 4 0 1 0 2 1 0 0 0 0 2 1 0 0 1 0 0 0 0	324	6
0	16	0 0 0 0 0 0 0 0 4 0 2 0 2 0 0 0 0 0 2 0 0 0 2 0 0 0 0	54	7
0	56	0 0 0 0 0 0 0 0 4 1 1 0 1 1 0 0 0 0 1 1 0 1 1 0 0 0 0	27	8
0	12	0 0 0 0 0 0 0 1 3 0 0 0 1 2 1 0 0 0 3 1 0 0 0 0 0 0 0	324	9
0	4	0 0 0 0 0 0 0 1 3 0 0 0 1 3 0 0 0 0 3 0 1 0 0 0 0 0 0	216	10
0	−12	0 0 0 0 0 0 0 1 3 0 0 0 2 1 1 0 0 0 2 2 0 0 0 0 0 0 0	648	11
0	4	0 0 0 0 0 0 0 1 3 0 0 1 0 3 0 0 0 0 3 0 0 1 0 0 0 0 0	648	12
0	−4	0 0 0 0 0 0 0 1 3 0 0 1 1 2 0 0 0 0 2 1 0 1 0 0 0 0 0	1296	13
0	−8	0 0 0 0 0 0 0 1 3 0 0 1 1 2 0 0 0 0 3 0 0 0 1 0 0 0 0	1296	14
0	16	0 0 0 0 0 0 0 1 3 0 0 1 2 1 0 0 0 0 2 1 0 0 1 0 0 0 0	648	15
0	12	0 0 0 0 0 0 0 1 3 0 1 0 0 2 1 0 0 0 3 0 0 1 0 0 0 0 0	648	16
0	−8	0 0 0 0 0 0 0 1 3 0 1 0 1 1 1 0 0 0 2 1 0 1 0 0 0 0 0	1296	17
0	−16	0 0 0 0 0 0 0 1 3 0 1 0 1 1 1 0 0 0 3 0 0 0 1 0 0 0 0	648	18
0	−4	0 0 0 0 0 0 0 1 3 0 1 0 1 2 0 0 0 0 2 0 1 1 0 0 0 0 0	1296	19
0	16	0 0 0 0 0 0 0 1 3 0 1 0 2 0 1 0 0 0 2 1 0 0 1 0 0 0 0	648	20
0	−12	0 0 0 0 0 0 0 1 3 0 1 1 0 2 0 0 0 0 2 0 0 2 0 0 0 0 0	1296	21
0	−8	0 0 0 0 0 0 0 1 3 0 1 1 1 1 0 0 0 0 1 1 0 2 0 0 0 0 0	648	22
0	32	0 0 0 0 0 0 0 1 3 0 1 1 1 1 0 0 0 0 2 0 0 1 1 0 0 0 0	1296	23
0	−32	0 0 0 0 0 0 0 1 3 0 1 1 2 0 0 0 0 0 2 0 0 0 2 0 0 0 0	648	24
0	−4	0 0 0 0 0 0 0 1 3 0 2 0 1 0 1 0 0 0 1 1 0 2 0 0 0 0 0	1296	25
0	16	0 0 0 0 0 0 0 1 3 0 2 0 1 0 1 0 0 0 2 0 0 1 1 0 0 0 0	1296	26
0	16	0 0 0 0 0 0 0 1 3 0 2 0 1 1 0 0 0 0 2 0 0 1 0 1 0 0 0	1296	27
0	−24	0 0 0 0 0 0 0 1 3 0 2 1 1 0 0 0 0 0 1 0 0 2 1 0 0 0 0	648	28
0	−8	0 0 0 0 0 0 0 1 3 0 3 0 1 0 0 0 0 0 1 0 0 2 0 1 0 0 0	648	29
0	32	0 0 0 0 0 0 0 1 3 1 0 0 1 1 1 0 0 0 1 2 0 1 0 0 0 0 0	648	30
0	−56	0 0 0 0 0 0 0 1 3 1 0 1 1 1 0 0 0 0 1 1 0 1 1 0 0 0 0	648	31
0	6	0 0 0 0 0 0 0 2 2 0 0 0 2 0 2 0 0 0 2 2 0 0 0 0 0 0 0	108	32
0	24	0 0 0 0 0 0 0 2 2 0 0 0 2 1 1 0 0 0 2 1 1 0 0 0 0 0 0	162	33
0	8	0 0 0 0 0 0 0 2 2 0 0 1 1 1 1 0 0 0 2 1 0 1 0 0 0 0 0	1296	34
0	16	0 0 0 0 0 0 0 2 2 0 0 1 1 1 1 0 0 0 3 0 0 0 1 0 0 0 0	648	35
0	4	0 0 0 0 0 0 0 2 2 0 0 1 1 2 0 0 0 0 2 0 1 1 0 0 0 0 0	1296	36
0	8	0 0 0 0 0 0 0 2 2 0 0 1 1 2 0 0 0 0 3 0 0 0 0 1 0 0 0	648	37
0	−16	0 0 0 0 0 0 0 2 2 0 0 1 2 0 1 0 0 0 2 1 0 0 1 0 0 0 0	648	38
0	−16	0 0 0 0 0 0 0 2 2 0 0 1 2 1 0 0 0 0 2 1 0 0 0 1 0 0 0	324	39
0	6	0 0 0 0 0 0 0 2 2 0 0 2 0 2 0 0 0 0 2 0 0 2 0 0 0 0 0	324	40
0	4	0 0 0 0 0 0 0 2 2 0 0 2 1 1 0 0 0 0 1 1 0 2 0 0 0 0 0	648	41
0	−16	0 0 0 0 0 0 0 2 2 0 0 2 1 1 0 0 0 0 2 0 0 1 1 0 0 0 0	1296	42
0	16	0 0 0 0 0 0 0 2 2 0 0 2 2 0 0 0 0 0 2 0 0 0 2 0 0 0 0	324	43
0	24	0 0 0 0 0 0 0 2 2 0 1 1 0 1 1 0 0 0 2 0 0 2 0 0 0 0 0	162	44
0	8	0 0 0 0 0 0 0 2 2 0 1 1 1 0 1 0 0 0 1 1 0 2 0 0 0 0 0	648	45
0	−32	0 0 0 0 0 0 0 2 2 0 1 1 1 0 1 0 0 0 2 0 0 1 1 0 0 0 0	1296	46
0	64	0 0 0 0 0 0 0 2 2 0 1 1 2 0 0 0 0 0 2 0 0 0 1 1 0 0 0	162	47
0	24	0 0 0 0 0 0 0 2 2 0 1 2 1 0 0 0 0 0 1 0 0 2 1 0 0 0 0	648	48
0	−16	0 0 0 0 0 0 0 2 2 1 0 0 1 0 2 0 0 0 1 2 0 1 0 0 0 0 0	324	49
0	−64	0 0 0 0 0 0 0 2 2 1 0 0 1 1 1 0 0 0 1 1 1 1 0 0 0 0 0	162	50
0	56	0 0 0 0 0 0 0 2 2 1 0 1 1 0 1 0 0 0 1 1 0 1 1 0 0 0 0	324	51
0	56	0 0 0 0 0 0 0 2 2 1 0 1 1 1 0 0 0 0 1 1 0 1 0 1 0 0 0	162	52
0	−12	0 0 0 0 0 0 1 1 2 0 0 0 1 2 1 0 0 0 2 1 1 0 0 0 0 0 0	108	53
0	28	0 0 0 0 0 0 1 1 2 0 0 1 0 2 1 0 0 0 2 1 0 1 0 0 0 0 0	1296	54
0	−4	0 0 0 0 0 0 1 1 2 0 0 1 0 2 1 0 0 0 3 0 0 0 1 0 0 0 0	1296	55
0	−8	0 0 0 0 0 0 1 1 2 0 0 1 0 3 0 0 0 0 3 0 0 0 0 1 0 0 0	324	56
0	−24	0 0 0 0 0 0 1 1 2 0 0 1 1 1 1 0 0 0 1 2 0 1 0 0 0 0 0	1296	57
0	8	0 0 0 0 0 0 1 1 2 0 0 1 1 2 0 0 0 0 2 1 0 0 0 1 0 0 0	324	58
0	−16	0 0 0 0 0 0 1 1 2 0 0 2 0 2 0 0 0 0 1 1 0 2 0 0 0 0 0	1296	59
0	4	0 0 0 0 0 0 1 1 2 0 0 2 0 2 0 0 0 0 2 0 0 1 1 0 0 0 0	648	60
0	56	0 0 0 0 0 0 1 1 2 0 0 2 1 1 0 0 0 0 1 1 0 1 1 0 0 0 0	324	61
0	8	0 0 0 0 0 0 1 1 2 0 1 0 0 1 2 0 0 0 2 1 0 1 0 0 0 0 0	1296	62
0	16	0 0 0 0 0 0 1 1 2 0 1 0 0 1 2 0 0 0 3 0 0 0 1 0 0 0 0	648	63
0	−32	0 0 0 0 0 0 1 1 2 0 1 0 0 2 1 0 0 0 2 0 1 1 0 0 0 0 0	648	64
0	−32	0 0 0 0 0 0 1 1 2 0 1 0 1 0 2 0 0 0 2 1 0 0 1 0 0 0 0	648	65
0	32	0 0 0 0 0 0 1 1 2 0 1 0 1 1 1 0 0 0 1 1 1 1 0 0 0 0 0	324	66
0	16	0 0 0 0 0 0 1 1 2 0 1 0 1 1 1 0 0 0 2 0 1 0 1 0 0 0 0	648	67
0	−24	0 0 0 0 0 0 1 1 2 0 1 1 0 1 1 0 0 0 1 1 0 2 0 0 0 0 0	1296	68
0	8	0 0 0 0 0 0 1 1 2 0 1 1 0 2 0 0 0 0 1 0 1 2 0 0 0 0 0	648	69
0	28	0 0 0 0 0 0 1 1 2 0 1 1 0 2 0 0 0 0 2 0 0 1 0 1 0 0 0	648	70
0	−8	0 0 0 0 0 0 1 1 2 0 1 1 1 0 1 0 0 0 1 1 0 1 1 0 0 0 0	648	71
0	48	0 0 0 0 0 0 1 1 2 0 1 1 1 0 1 0 0 0 2 0 0 0 2 0 0 0 0	648	72

TABLE 7. The basis invariants in degree 12: part 1

I_{12}	I'_{12}	MINIMAL REPRESENTATIVE	ORBIT SIZE	#
0	72	0 0 0 0 0 0 1 1 2 0 1 1 1 1 0 0 0 0 1 1 0 1 0 1 0 0 0	324	73
0	−64	0 0 0 0 0 0 1 1 2 0 1 1 1 1 0 0 0 0 2 0 0 0 1 1 0 0 0	648	74
0	−12	0 0 0 0 0 1 0 1 2 0 0 1 0 2 0 1 0 0 2 1 0 1 0 0 0 0 0	324	75
0	16	0 0 0 0 0 1 0 1 2 0 0 1 0 2 0 1 0 0 3 0 0 0 1 0 0 0 0	648	76
1	13	0 0 0 0 0 1 0 1 2 0 0 1 1 1 0 0 1 0 2 1 0 1 0 0 0 0 0	648	77
−1	−5	0 0 0 0 0 1 0 1 2 0 0 1 1 1 0 0 1 0 3 0 0 0 1 0 0 0 0	1296	78
−1	19	0 0 0 0 0 1 0 1 2 0 0 1 1 1 0 1 0 0 1 2 0 1 0 0 0 0 0	1296	79
1	−27	0 0 0 0 0 1 0 1 2 0 0 1 1 1 0 1 0 0 2 1 0 0 1 0 0 0 0	1296	80
−1	11	0 0 0 0 0 1 0 1 2 0 0 1 1 2 0 0 0 0 2 1 0 0 0 0 1 0 0	1296	81
1	13	0 0 0 0 0 1 0 1 2 0 0 1 1 2 0 0 0 0 3 0 0 0 0 0 0 1 0	1296	82
−1	−9	0 0 0 0 0 1 0 1 2 0 0 1 2 0 0 0 1 0 1 2 0 1 0 0 0 0 0	1296	83
1	5	0 0 0 0 0 1 0 1 2 0 0 1 2 0 0 0 1 0 2 1 0 0 1 0 0 0 0	1296	84
1	−7	0 0 0 0 0 1 0 1 2 0 0 1 2 0 0 1 0 0 0 3 0 1 0 0 0 0 0	648	85
2	2	0 0 0 0 0 1 0 1 2 0 0 1 2 1 0 0 0 0 1 2 0 0 0 0 1 0 0	1296	86
−2	−26	0 0 0 0 0 1 0 1 2 0 0 1 2 1 0 0 0 0 2 1 0 0 0 0 0 1 0	1296	87
−1	−5	0 0 0 0 0 1 0 1 2 0 0 1 3 0 0 0 0 0 0 3 0 0 0 0 1 0 0	648	88
1	13	0 0 0 0 0 1 0 1 2 0 0 1 3 0 0 0 0 0 1 2 0 0 0 0 0 1 0	648	89
−2	−18	0 0 0 0 0 1 0 1 2 0 1 0 1 0 1 0 1 0 2 1 0 1 0 0 0 0 0	324	90
2	26	0 0 0 0 0 1 0 1 2 0 1 0 1 0 1 0 1 0 3 0 0 0 1 0 0 0 0	648	91
1	13	0 0 0 0 0 1 0 1 2 0 1 0 1 0 1 1 0 0 1 2 0 1 0 0 0 0 0	1296	92
−1	−21	0 0 0 0 0 1 0 1 2 0 1 0 1 0 1 1 0 0 2 1 0 0 1 0 0 0 0	1296	93
0	16	0 0 0 0 0 1 0 1 2 0 1 0 1 1 0 0 0 1 2 1 0 1 0 0 0 0 0	648	94
0	32	0 0 0 0 0 1 0 1 2 0 1 0 1 1 0 0 0 1 3 0 0 0 1 0 0 0 0	216	95
1	13	0 0 0 0 0 1 0 1 2 0 1 0 1 1 0 0 1 0 2 0 1 1 0 0 0 0 0	1296	96
−1	−5	0 0 0 0 0 1 0 1 2 0 1 0 1 1 0 0 1 0 3 0 0 0 0 1 0 0 0	1296	97
0	−8	0 0 0 0 0 1 0 1 2 0 1 0 1 1 0 1 0 0 1 1 1 1 0 0 0 0 0	1296	98
1	−27	0 0 0 0 0 1 0 1 2 0 1 0 1 1 0 1 0 0 2 0 1 0 1 0 0 0 0	1296	99
−1	−21	0 0 0 0 0 1 0 1 2 0 1 0 1 1 0 1 0 0 2 1 0 0 0 1 0 0 0	1296	100
0	−12	0 0 0 0 0 1 0 1 2 0 1 0 1 2 0 0 0 0 2 0 1 0 0 0 1 0 0	216	101
0	16	0 0 0 0 0 1 0 1 2 0 1 0 1 2 0 0 0 0 3 0 0 0 0 0 0 0 1	648	102
0	8	0 0 0 0 0 1 0 1 2 0 1 0 2 0 0 0 0 1 1 2 0 1 0 0 0 0 0	1296	103
0	−32	0 0 0 0 0 1 0 1 2 0 1 0 2 0 0 0 0 1 2 1 0 0 1 0 0 0 0	648	104
1	13	0 0 0 0 0 1 0 1 2 0 1 0 2 0 0 0 1 0 1 1 1 1 0 0 0 0 0	1296	105
−2	−26	0 0 0 0 0 1 0 1 2 0 1 0 2 0 0 0 1 0 2 0 1 0 1 0 0 0 0	1296	106
1	5	0 0 0 0 0 1 0 1 2 0 1 0 2 0 0 0 1 0 2 1 0 0 0 1 0 0 0	1296	107
−1	−29	0 0 0 0 0 1 0 1 2 0 1 0 2 0 0 1 0 0 0 2 1 1 0 0 0 0 0	216	108
1	53	0 0 0 0 0 1 0 1 2 0 1 0 2 0 0 1 0 0 1 1 1 0 1 0 0 0 0	1296	109
0	8	0 0 0 0 0 1 0 1 2 0 1 0 2 0 0 1 0 0 1 2 0 0 0 1 0 0 0	1296	110
−1	−9	0 0 0 0 0 1 0 1 2 0 1 0 2 0 1 0 0 0 1 2 0 0 0 0 1 0 0	1296	111
1	5	0 0 0 0 0 1 0 1 2 0 1 0 2 0 1 0 0 0 2 1 0 0 0 0 0 1 0	1296	112
−1	19	0 0 0 0 0 1 0 1 2 0 1 0 2 1 0 0 0 0 1 1 1 0 0 0 1 0 0	1296	113
1	5	0 0 0 0 0 1 0 1 2 0 1 0 2 1 0 0 0 0 2 0 1 0 0 0 0 1 0	1296	114
0	−32	0 0 0 0 0 1 0 1 2 0 1 0 2 1 0 0 0 0 2 1 0 0 0 0 0 0 1	648	115
−1	−5	0 0 0 0 0 1 0 1 2 0 1 0 3 0 0 0 0 0 1 1 1 0 0 0 0 1 0	1296	116
−2	14	0 0 0 0 0 1 0 1 2 0 1 1 1 0 0 1 0 0 0 2 0 2 0 0 0 0 0	648	117
4	−60	0 0 0 0 0 1 0 1 2 0 1 1 1 0 0 1 0 0 1 1 0 1 1 0 0 0 0	648	118
−2	54	0 0 0 0 0 1 0 1 2 0 1 1 1 0 0 1 0 0 2 0 0 0 2 0 0 0 0	648	119
2	22	0 0 0 0 0 1 0 1 2 0 1 1 2 0 0 0 0 0 0 2 0 1 0 0 1 0 0	648	120
−3	−15	0 0 0 0 0 1 0 1 2 0 1 1 2 0 0 0 0 0 1 1 0 0 1 0 1 0 0	1296	121
−1	−21	0 0 0 0 0 1 0 1 2 0 1 1 2 0 0 0 0 0 1 1 0 1 0 0 0 1 0	1296	122
2	42	0 0 0 0 0 1 0 1 2 0 1 1 2 0 0 0 0 0 2 0 0 0 1 0 0 1 0	1296	123
2	−38	0 0 0 0 0 1 0 1 2 0 2 0 2 0 0 0 0 0 1 0 1 0 1 0 1 0 0	648	124
1	5	0 0 0 0 0 1 0 1 2 0 2 0 2 0 0 0 0 0 1 0 1 1 0 0 0 1 0	1296	125
−2	−10	0 0 0 0 0 1 0 1 2 0 2 0 2 0 0 0 0 0 2 0 0 0 0 1 0 1 0	648	126
0	64	0 0 0 0 0 1 0 1 2 0 2 0 2 0 0 0 0 0 2 0 0 0 1 0 0 0 1	216	127
−1	−21	0 0 0 0 0 1 0 1 2 1 0 0 0 1 1 1 0 0 1 2 0 1 0 0 0 0 0	1296	128
1	13	0 0 0 0 0 1 0 1 2 1 0 0 0 1 1 1 0 0 2 1 0 0 1 0 0 0 0	1296	129
0	48	0 0 0 0 0 1 0 1 2 1 0 0 0 2 0 0 0 1 2 1 0 1 0 0 0 0 0	108	130
1	−27	0 0 0 0 0 1 0 1 2 1 0 0 0 2 0 1 0 0 1 1 1 1 0 0 0 0 0	1296	131
−1	−9	0 0 0 0 0 1 0 1 2 1 0 0 0 2 0 1 0 0 2 1 0 0 0 1 0 0 0	648	132
2	10	0 0 0 0 0 1 0 1 2 1 0 0 1 0 1 0 1 0 1 2 0 1 0 0 0 0 0	1296	133
−2	−18	0 0 0 0 0 1 0 1 2 1 0 0 1 0 1 0 1 0 2 1 0 0 1 0 0 0 0	648	134
0	−64	0 0 0 0 0 1 0 1 2 1 0 0 1 1 0 0 0 1 1 2 0 1 0 0 0 0 0	648	135
−2	−42	0 0 0 0 0 1 0 1 2 1 0 0 1 1 0 0 1 0 1 1 1 1 0 0 0 0 0	1296	136
−2	−2	0 0 0 0 0 1 0 1 2 1 0 0 1 1 0 1 0 0 1 1 1 0 1 0 0 0 0	648	137
1	13	0 0 0 0 0 1 0 1 2 1 0 0 1 1 0 1 0 0 1 2 0 0 0 1 0 0 0	1296	138
−1	−21	0 0 0 0 0 1 0 1 2 1 0 0 1 1 1 0 0 0 1 2 0 0 0 0 1 0 0	1296	139
1	13	0 0 0 0 0 1 0 1 2 1 0 0 1 1 1 0 0 0 2 1 0 0 0 0 0 1 0	648	140
1	−27	0 0 0 0 0 1 0 1 2 1 0 0 1 2 0 0 0 0 1 1 1 0 0 0 1 0 0	1296	141
−1	−9	0 0 0 0 0 1 0 1 2 1 0 0 2 0 0 0 1 0 1 2 0 0 0 1 0 0 0	648	142
−1	−9	0 0 0 0 0 1 0 1 2 1 0 0 2 0 1 0 0 0 1 2 0 0 0 0 0 1 0	216	143
−2	78	0 0 0 0 0 1 0 1 2 1 0 1 0 1 0 1 0 0 1 1 0 1 1 0 0 0 0	648	144

TABLE 8. The basis invariants in degree 12: part 2

I_{12}	I'_{12}	MINIMAL REPRESENTATIVE	ORBIT SIZE	#
0	−32	0 0 0 0 0 1 0 1 2 1 0 1 0 1 0 1 0 0 2 0 0 0 2 0 0 0 0	648	145
−2	−2	0 0 0 0 0 1 0 1 2 1 0 1 1 0 0 0 1 0 1 1 0 1 1 0 0 0 0	1296	146
1	5	0 0 0 0 0 1 0 1 2 1 0 1 1 0 0 0 1 0 2 0 0 0 2 0 0 0 0	1296	147
0	32	0 0 0 0 0 1 0 1 2 1 0 1 1 1 0 0 0 0 1 1 0 0 1 0 1 0 0	648	148
4	60	0 0 0 0 0 1 0 1 2 1 0 1 1 1 0 0 0 0 1 1 0 1 0 0 0 1 0	1296	149
−1	−21	0 0 0 0 0 1 0 1 2 1 0 1 1 1 0 0 0 0 2 0 0 0 1 0 0 1 0	1296	150
0	112	0 0 0 0 0 1 0 1 2 1 1 0 1 0 0 0 0 1 1 1 0 1 1 0 0 0 0	216	151
−2	−2	0 0 0 0 0 1 0 1 2 1 1 0 1 0 0 0 1 0 1 1 0 1 0 1 0 0 0	648	152
−1	−21	0 0 0 0 0 1 0 1 2 1 1 0 1 0 0 0 1 0 2 0 0 0 1 1 0 0 0	1296	153
2	10	0 0 0 0 0 1 0 1 2 1 1 0 1 1 0 0 0 0 2 0 0 0 0 1 0 1 0	648	154
1	−11	0 0 0 0 0 1 0 2 1 0 0 1 0 1 1 1 0 0 3 0 0 0 1 0 0 0 0	1296	155
−1	−21	0 0 0 0 0 1 0 2 1 0 0 1 0 2 0 1 0 0 3 0 0 0 0 1 0 0 0	1296	156
0	−24	0 0 0 0 0 1 0 2 1 0 0 1 1 0 1 1 0 0 1 2 0 1 0 0 0 0 0	648	157
0	16	0 0 0 0 0 1 0 2 1 0 0 1 1 0 1 1 0 0 2 1 0 0 1 0 0 0 0	1296	158
1	−11	0 0 0 0 0 1 0 2 1 0 0 1 1 1 0 0 0 1 3 0 0 0 1 0 0 0 0	1296	159
2	−14	0 0 0 0 0 1 0 2 1 0 0 1 1 1 0 1 0 0 1 1 1 1 0 0 0 0 0	1296	160
−2	22	0 0 0 0 0 1 0 2 1 0 0 1 1 1 0 1 0 0 2 0 1 0 1 0 0 0 0	648	161
0	16	0 0 0 0 0 1 0 2 1 0 0 1 1 1 0 1 0 0 2 1 0 0 0 1 0 0 0	1296	162
−1	−21	0 0 0 0 0 1 0 2 1 0 0 1 1 2 0 0 0 0 3 0 0 0 0 0 0 0 1	1296	163
1	5	0 0 0 0 0 1 0 2 1 0 0 1 2 0 0 0 0 1 1 2 0 1 0 0 0 0 0	1296	164
−1	11	0 0 0 0 0 1 0 2 1 0 0 1 2 0 0 0 0 1 2 1 0 0 1 0 0 0 0	1296	165
1	−27	0 0 0 0 0 1 0 2 1 0 0 1 2 0 0 1 0 0 1 1 1 0 1 0 0 0 0	1296	166
1	5	0 0 0 0 0 1 0 2 1 0 0 1 2 0 0 1 0 0 1 2 0 0 0 1 0 0 0	1296	167
−1	11	0 0 0 0 0 1 0 2 1 0 0 1 2 0 1 0 0 0 1 2 0 0 0 0 1 0 0	1296	168
−3	−15	0 0 0 0 0 1 0 2 1 0 0 1 2 1 0 0 0 0 1 1 1 0 0 0 1 0 0	1296	169
2	42	0 0 0 0 0 1 0 2 1 0 0 1 2 1 0 0 0 0 2 1 0 0 0 0 0 0 1	648	170
2	10	0 0 0 0 0 1 0 2 1 0 0 1 3 0 0 0 0 0 0 2 1 0 0 0 1 0 0	648	171
1	−15	0 0 0 0 0 1 0 2 1 0 0 2 0 1 0 1 0 0 1 1 0 2 0 0 0 0 0	648	172
−1	11	0 0 0 0 0 1 0 2 1 0 0 2 0 1 0 1 0 0 2 0 0 1 1 0 0 0 0	1296	173
2	−2	0 0 0 0 0 1 0 2 1 0 0 2 1 0 0 1 0 0 0 2 0 2 0 0 0 0 0	648	174
−4	28	0 0 0 0 0 1 0 2 1 0 0 2 1 0 0 1 0 0 1 1 0 1 1 0 0 0 0	1296	175
2	−22	0 0 0 0 0 1 0 2 1 0 0 2 1 0 0 1 0 0 2 0 0 0 2 0 0 0 0	1296	176
0	16	0 0 0 0 0 1 0 2 1 0 0 2 1 1 0 0 0 0 1 1 0 1 0 0 1 0 0	1296	177
−1	11	0 0 0 0 0 1 0 2 1 0 0 2 1 1 0 0 0 0 2 0 0 0 1 0 1 0 0	1296	178
−2	−10	0 0 0 0 0 1 0 2 1 0 0 2 2 0 0 0 0 0 0 2 0 1 0 0 1 0 0	648	179
3	−1	0 0 0 0 0 1 0 2 1 0 0 2 2 0 0 0 0 0 1 1 0 0 1 0 1 0 0	1296	180
1	13	0 0 0 0 0 1 0 2 1 0 1 0 0 0 2 1 0 0 3 0 0 0 1 0 0 0 0	648	181
−1	−5	0 0 0 0 0 1 0 2 1 0 1 0 0 1 1 1 0 0 3 0 0 0 0 1 0 0 0	1296	182
−2	−42	0 0 0 0 0 1 0 2 1 0 1 0 1 0 1 0 0 1 3 0 0 0 1 0 0 0 0	648	183
−2	−2	0 0 0 0 0 1 0 2 1 0 1 0 1 0 1 1 0 0 1 1 1 1 0 0 0 0 0	1296	184
0	16	0 0 0 0 0 1 0 2 1 0 1 0 1 0 1 1 0 0 2 0 1 0 1 0 0 0 0	1296	185
2	10	0 0 0 0 0 1 0 2 1 0 1 0 1 0 1 1 0 0 2 1 0 0 0 1 0 0 0	1296	186
0	16	0 0 0 0 0 1 0 2 1 0 1 0 1 1 0 1 0 0 2 0 1 0 0 1 0 0 0	648	187
1	−11	0 0 0 0 0 1 0 2 1 0 1 0 1 1 1 0 0 0 3 0 0 0 0 0 0 0 1	1296	188
−1	−21	0 0 0 0 0 1 0 2 1 0 1 0 2 0 0 0 0 1 1 1 1 1 0 0 0 0 0	1296	189
2	42	0 0 0 0 0 1 0 2 1 0 1 0 2 0 0 0 0 1 2 0 1 0 1 0 0 0 0	648	190
−1	−29	0 0 0 0 0 1 0 2 1 0 1 0 2 0 0 1 0 0 1 0 2 0 1 0 0 0 0	1296	191
−1	−21	0 0 0 0 0 1 0 2 1 0 1 0 2 0 0 1 0 0 1 1 1 0 0 1 0 0 0	1296	192
3	7	0 0 0 0 0 1 0 2 1 0 1 0 2 0 1 0 0 0 1 1 1 0 0 0 1 0 0	648	193
1	−15	0 0 0 0 0 1 0 2 1 0 1 0 2 1 0 0 0 0 1 0 2 0 0 0 1 0 0	432	194
−2	2	0 0 0 0 0 1 0 2 1 0 1 0 3 0 0 0 0 0 0 1 2 0 0 0 1 0 0	648	195
3	7	0 0 0 0 0 1 0 2 1 0 1 1 0 0 1 1 0 0 1 1 0 2 0 0 0 0 0	648	196
−3	−15	0 0 0 0 0 1 0 2 1 0 1 1 0 0 1 1 0 0 2 0 0 1 1 0 0 0 0	1296	197
3	47	0 0 0 0 0 1 0 2 1 0 1 1 0 1 0 1 0 0 2 0 0 1 0 1 0 0 0	648	198
1	53	0 0 0 0 0 1 0 2 1 0 1 1 1 0 0 0 0 1 2 0 0 1 1 0 0 0 0	1296	199
0	−24	0 0 0 0 0 1 0 2 1 0 1 1 1 0 0 1 0 0 0 1 1 2 0 0 0 0 0	648	200
2	26	0 0 0 0 0 1 0 2 1 0 1 1 1 0 0 1 0 0 1 0 1 1 1 0 0 0 0	1296	201
−2	38	0 0 0 0 0 1 0 2 1 0 1 1 1 0 0 1 0 0 1 1 0 1 0 1 0 0 0	1296	202
0	−64	0 0 0 0 0 1 0 2 1 0 1 1 1 0 0 1 0 0 2 0 0 0 1 1 0 0 0	1296	203
−2	−2	0 0 0 0 0 1 0 2 1 0 1 1 1 0 1 0 0 0 1 1 0 1 0 0 1 0 0	1296	204
3	47	0 0 0 0 0 1 0 2 1 0 1 1 1 0 1 0 0 0 2 0 0 0 1 0 1 0 0	1296	205
2	−14	0 0 0 0 0 1 0 2 1 0 1 1 1 1 0 0 0 0 1 0 1 1 0 0 1 0 0	1296	206
−3	−15	0 0 0 0 0 1 0 2 1 0 1 1 1 1 0 0 0 0 2 0 0 0 0 1 1 0 0	1296	207
1	53	0 0 0 0 0 1 0 2 1 0 1 1 1 1 0 0 0 0 2 0 0 1 0 0 0 0 1	1296	208
−1	59	0 0 0 0 0 1 0 2 1 0 1 1 2 0 0 0 0 0 1 0 1 0 1 0 1 0 0	1296	209
1	−27	0 0 0 0 0 1 0 2 1 0 1 1 2 0 0 0 0 0 1 1 0 0 0 1 1 0 0	1296	210
−2	−106	0 0 0 0 0 1 0 2 1 0 1 1 2 0 0 0 0 0 2 0 0 0 1 0 0 0 1	648	211
−2	−58	0 0 0 0 0 1 0 2 1 0 1 2 1 0 0 0 0 0 1 0 0 1 1 0 1 0 0	1296	212
1	5	0 0 0 0 0 1 0 2 1 0 2 0 0 0 1 1 0 0 2 0 0 1 0 1 0 0 0	648	213
−1	11	0 0 0 0 0 1 0 2 1 0 2 0 1 0 0 0 0 1 2 0 0 1 0 1 0 0 0	648	214
4	4	0 0 0 0 0 1 0 2 1 0 2 0 1 0 0 1 0 0 1 0 1 1 0 1 0 0 0	1296	215
−2	−10	0 0 0 0 0 1 0 2 1 0 2 0 1 0 0 1 0 0 2 0 0 0 0 2 0 0 0	648	216

Table 9. The basis invariants in degree 12: part 3

I_{12}	I'_{12}	MINIMAL REPRESENTATIVE	ORBIT SIZE	#
0	−24	0 0 0 0 0 1 0 2 1 0 2 0 1 0 1 0 0 0 1 0 1 1 0 0 1 0 0	1296	217
1	5	0 0 0 0 0 1 0 2 1 0 2 0 1 0 1 0 0 0 2 0 0 0 0 1 1 0 0	1296	218
−1	11	0 0 0 0 0 1 0 2 1 0 2 0 1 0 1 0 0 0 2 0 0 1 0 0 0 0 1	1296	219
−3	17	0 0 0 0 0 1 0 2 1 0 2 0 2 0 0 0 0 0 1 0 1 0 0 1 1 0 0	1296	220
2	−22	0 0 0 0 0 1 0 2 1 0 2 0 2 0 0 0 0 0 2 0 0 0 0 1 0 0 1	648	221
2	10	0 0 0 0 0 1 0 2 1 0 2 1 1 0 0 0 0 0 1 0 0 1 0 1 1 0 0	1296	222
0	32	0 0 0 0 0 1 0 2 1 1 0 0 0 1 1 1 0 0 1 1 1 1 0 0 0 0 0	1296	223
−2	22	0 0 0 0 0 1 0 2 1 1 0 0 1 0 1 0 0 1 1 2 0 1 0 0 0 0 0	324	224
0	−8	0 0 0 0 0 1 0 2 1 1 0 0 1 0 1 1 0 0 1 1 1 0 1 0 0 0 0	1296	225
−2	−18	0 0 0 0 0 1 0 2 1 1 0 0 1 0 1 1 0 0 1 2 0 0 0 1 0 0 0	648	226
2	106	0 0 0 0 0 1 0 2 1 1 0 0 1 1 0 0 0 1 1 1 1 1 0 0 0 0 0	648	227
0	−8	0 0 0 0 0 1 0 2 1 1 0 0 1 1 0 1 0 0 1 1 1 0 0 1 0 0 0	1296	228
0	32	0 0 0 0 0 1 0 2 1 1 0 0 1 1 1 0 0 0 1 1 1 0 0 0 1 0 0	1296	229
4	−20	0 0 0 0 0 1 0 2 1 1 0 1 0 0 1 1 0 0 1 1 0 1 1 0 0 0 0	1296	230
1	53	0 0 0 0 0 1 0 2 1 1 0 1 0 1 0 0 0 1 1 1 0 2 0 0 0 0 0	648	231
2	−54	0 0 0 0 0 1 0 2 1 1 0 1 0 1 0 1 0 0 1 0 1 1 1 0 0 0 0	1296	232
−2	−82	0 0 0 0 0 1 0 2 1 1 0 1 0 1 0 1 0 0 1 1 0 1 0 1 0 0 0	1296	233
0	32	0 0 0 0 0 1 0 2 1 1 0 1 0 1 1 0 0 0 1 1 0 1 0 0 1 0 0	1296	234
−2	22	0 0 0 0 0 1 0 2 1 1 0 1 0 2 0 0 0 0 1 0 1 1 0 0 1 0 0	1296	235
2	−54	0 0 0 0 0 1 0 2 1 1 0 1 1 0 0 0 0 1 1 1 0 1 1 0 0 0 0	1296	236
0	32	0 0 0 0 0 1 0 2 1 1 0 1 1 0 0 1 0 0 1 1 0 0 1 1 0 0 0	1296	237
−4	−36	0 0 0 0 0 1 0 2 1 1 0 1 1 0 1 0 0 0 1 1 0 0 1 0 1 0 0	1296	238
2	26	0 0 0 0 0 1 0 2 1 1 0 1 1 1 0 0 0 0 1 1 0 0 0 1 1 0 0	1296	239
−4	−116	0 0 0 0 0 1 0 2 1 1 0 1 1 1 0 0 0 0 1 1 0 1 0 0 0 0 1	648	240
−2	22	0 0 0 0 0 1 0 2 1 1 1 0 0 0 1 0 0 1 1 1 0 2 0 0 0 0 0	648	241
−2	−2	0 0 0 0 0 1 0 2 1 1 1 0 0 0 1 1 0 0 1 1 0 1 0 1 0 0 0	1296	242
2	10	0 0 0 0 0 1 0 2 1 1 1 0 0 0 2 0 0 0 1 1 0 1 0 0 1 0 0	648	243
1	21	0 0 0 0 0 1 0 3 0 0 0 1 0 1 1 1 0 0 3 0 0 0 0 1 0 0 0	648	244
0	24	0 0 0 0 0 1 0 3 0 0 0 1 1 0 1 1 0 0 1 1 1 1 0 0 0 0 0	648	245
−1	−5	0 0 0 0 0 1 0 3 0 0 0 1 1 0 1 1 0 0 2 1 0 0 0 1 0 0 0	1296	246
−2	−10	0 0 0 0 0 1 0 3 0 0 0 1 1 1 0 0 0 1 3 0 0 0 0 1 0 0 0	108	247
2	10	0 0 0 0 0 1 0 3 0 0 0 1 2 0 0 0 0 1 2 1 0 0 0 1 0 0 0	648	248
3	−17	0 0 0 0 0 1 0 3 0 0 0 2 1 0 0 1 0 0 1 1 0 1 0 1 0 0 0	432	249
−2	22	0 0 0 0 0 1 0 3 0 0 0 2 1 0 0 1 0 0 2 0 0 0 1 1 0 0 0	648	250
2	42	0 0 0 0 0 1 0 3 0 0 0 2 2 0 0 0 0 0 2 0 0 0 1 0 0 0 1	216	251
1	1	0 0 0 0 0 1 0 3 0 0 0 3 0 0 0 1 0 0 0 1 0 3 0 0 0 0 0	72	252
−4	−4	0 0 0 0 0 1 0 3 0 0 1 1 1 0 0 1 0 0 1 0 1 1 0 1 0 0 0	648	253
2	−22	0 0 0 0 0 1 0 3 0 1 0 0 1 0 1 0 0 1 1 1 1 1 0 0 0 0 0	648	254
2	18	0 0 0 0 0 1 0 3 0 1 0 0 1 0 1 1 0 0 1 1 1 0 0 1 0 0 0	216	255
2	58	0 0 0 0 0 1 0 3 0 1 0 1 0 1 0 1 0 0 1 0 1 1 0 1 0 0 0	648	256
−4	4	0 0 0 0 0 1 1 1 1 0 0 1 0 2 0 1 0 0 1 1 1 1 0 0 0 0 0	324	257
3	47	0 0 0 0 0 1 1 1 1 0 0 1 0 2 0 1 0 0 2 1 0 0 0 1 0 0 0	1296	258
0	24	0 0 0 0 0 1 1 1 1 0 0 1 1 1 0 0 1 0 1 1 1 1 0 0 0 0 0	648	259
−2	−42	0 0 0 0 0 1 1 1 1 0 0 1 1 1 0 0 1 0 2 1 0 0 0 1 0 0 0	1296	260
4	20	0 0 0 0 0 1 1 1 1 0 0 1 1 2 0 0 0 0 1 1 1 0 0 0 1 0 0	648	261
−1	−21	0 0 0 0 0 1 1 1 1 0 0 1 1 2 0 0 0 0 2 1 0 0 0 0 0 0 1	648	262
−3	17	0 0 0 0 0 1 1 1 1 0 0 2 0 1 0 1 0 0 0 2 0 2 0 0 0 0 0	648	263
4	−20	0 0 0 0 0 1 1 1 1 0 0 2 0 1 0 1 0 0 1 1 0 1 1 0 0 0 0	1296	264
−1	11	0 0 0 0 0 1 1 1 1 0 0 2 0 1 0 1 0 0 2 0 0 0 2 0 0 0 0	1296	265
0	16	0 0 0 0 0 1 1 1 1 0 0 2 0 2 0 0 0 0 1 1 0 1 0 0 1 0 0	648	266
4	4	0 0 0 0 0 1 1 1 1 0 0 2 1 1 0 0 0 0 0 2 0 1 0 0 1 0 0	1296	267
−4	−36	0 0 0 0 0 1 1 1 1 0 0 2 1 1 0 0 0 0 1 1 0 0 1 0 1 0 0	1296	268
0	32	0 0 0 0 0 1 1 1 1 0 1 0 0 1 1 1 0 0 2 0 1 0 1 0 0 0 0	1296	269
2	18	0 0 0 0 0 1 1 1 1 0 1 0 1 0 1 0 1 0 1 1 1 1 0 0 0 0 0	648	270
2	−22	0 0 0 0 0 1 1 1 1 0 1 0 1 0 1 1 0 0 1 1 1 0 1 0 0 0 0	648	271
−2	−42	0 0 0 0 0 1 1 1 1 0 1 0 1 1 0 0 0 1 2 0 1 0 1 0 0 0 0	648	272
−4	−84	0 0 0 0 0 1 1 1 1 0 1 0 1 1 1 0 0 0 1 1 1 0 0 0 1 0 0	216	273
−2	−42	0 0 0 0 0 1 1 1 1 0 1 0 2 0 0 0 0 1 1 1 1 0 1 0 0 0 0	324	274
−2	−2	0 0 0 0 0 1 1 1 1 0 1 0 2 0 0 0 1 0 1 1 1 0 0 1 0 0 0	324	275
−4	36	0 0 0 0 0 1 1 1 1 0 1 1 0 0 1 1 0 0 1 1 0 1 1 0 0 0 0	1296	276
2	58	0 0 0 0 0 1 1 1 1 0 1 1 0 1 0 1 0 0 1 0 1 1 1 0 0 0 0	648	277
−2	−50	0 0 0 0 0 1 1 1 1 0 1 1 0 1 0 1 0 0 1 1 0 1 0 1 0 0 0	1296	278
2	18	0 0 0 0 0 1 1 1 1 0 1 1 0 1 1 0 0 0 1 1 0 1 0 0 1 0 0	648	279
0	32	0 0 0 0 0 1 1 1 1 0 1 1 0 1 1 0 0 0 2 0 0 1 0 0 0 1 0	1296	280
−2	−10	0 0 0 0 0 1 1 1 1 0 1 1 1 0 0 0 0 1 1 1 0 1 1 0 0 0 0	1296	281
0	−64	0 0 0 0 0 1 1 1 1 0 1 1 1 0 0 0 0 1 2 0 0 0 2 0 0 0 0	1296	282
0	−56	0 0 0 0 0 1 1 1 1 0 1 1 1 0 0 0 1 0 1 1 0 1 0 1 0 0 0	1296	283
2	106	0 0 0 0 0 1 1 1 1 0 1 1 1 0 0 0 1 0 2 0 0 0 1 1 0 0 0	648	284
0	−48	0 0 0 0 0 1 1 1 1 0 1 1 1 0 0 1 0 0 1 0 1 0 2 0 0 0 0	1296	285
2	58	0 0 0 0 0 1 1 1 1 0 1 1 1 0 0 1 0 0 1 1 0 0 1 1 0 0 0	1296	286
6	46	0 0 0 0 0 1 1 1 1 0 1 1 1 0 1 0 0 0 1 1 0 1 0 0 0 1 0	648	287
−4	−36	0 0 0 0 0 1 1 1 1 0 1 1 1 0 1 0 0 0 2 0 0 0 1 0 0 1 0	1296	288

Table 10. The basis invariants in degree 12: part 4

I_{12}	I'_{12}	MINIMAL REPRESENTATIVE	ORBIT SIZE	#
0	−96	0 0 0 0 0 1 1 1 1 0 1 1 1 1 0 0 0 0 1 1 0 1 0 0 0 0 1	648	289
2	106	0 0 0 0 0 1 1 1 1 0 1 1 1 1 0 0 0 0 2 0 0 0 1 0 0 0 1	1296	290
4	−108	0 0 0 0 0 1 1 1 1 1 1 0 0 0 1 0 0 1 1 1 0 1 1 0 0 0 0	324	291
2	10	0 0 0 0 0 1 1 2 0 0 0 1 1 1 0 0 0 1 2 1 0 0 0 1 0 0 0	324	292
−4	−20	0 0 0 0 0 1 1 2 0 0 0 1 2 0 0 0 0 1 1 2 0 0 0 1 0 0 0	324	293
1	21	0 0 0 0 0 1 1 2 0 0 0 2 0 2 0 0 0 0 2 0 0 1 0 0 0 0 1	1296	294
−1	−13	0 0 0 0 0 1 1 2 0 0 0 2 1 0 0 0 0 1 0 2 0 2 0 0 0 0 0	432	295
0	16	0 0 0 0 0 1 1 2 0 0 0 2 1 0 0 0 0 1 1 1 0 1 1 0 0 0 0	1296	296
4	84	0 0 0 0 0 1 1 2 0 0 0 2 1 1 0 0 0 0 1 1 0 1 0 0 0 0 1	648	297
−3	−63	0 0 0 0 0 1 1 2 0 0 0 2 1 1 0 0 0 0 2 0 0 0 1 0 0 0 1	1296	298
2	10	0 0 0 0 0 1 1 2 0 0 1 0 1 0 1 0 0 1 2 0 1 0 1 0 0 0 0	648	299
−2	−26	0 0 0 0 0 1 1 2 0 0 1 0 2 0 0 0 0 1 1 0 2 0 1 0 0 0 0	324	300
0	−48	0 0 0 0 0 1 1 2 0 0 1 1 1 0 0 0 0 1 1 0 1 1 1 0 0 0 0	1296	301
2	26	0 0 0 0 0 1 1 2 0 0 1 1 1 0 0 0 0 1 1 1 0 1 0 1 0 0 0	1296	302
−2	22	0 0 0 0 0 1 1 2 0 0 1 1 1 0 0 0 0 1 2 0 0 0 1 1 0 0 0	648	303
−4	−36	0 0 0 0 0 1 1 2 0 0 1 1 1 0 1 0 0 0 1 1 0 1 0 0 0 0 1	1296	304
1	53	0 0 0 0 0 1 1 2 0 0 1 1 1 0 1 0 0 0 2 0 0 0 1 0 0 0 1	1296	305
3	−33	0 0 0 0 0 1 1 2 0 0 1 1 1 1 0 0 0 0 2 0 0 0 0 1 0 0 1	1296	306
−2	−58	0 0 0 0 0 1 1 2 0 0 2 0 1 0 0 0 0 1 1 0 1 1 0 1 0 0 0	1296	307
4	4	0 0 0 0 0 1 1 2 0 0 2 0 1 0 1 0 0 0 1 0 1 1 0 0 0 0 1	648	308
−2	22	0 0 0 0 0 1 1 2 0 0 2 1 1 0 0 0 0 0 1 0 0 1 0 1 0 0 1	324	309
2	26	0 0 0 0 0 1 1 2 0 1 0 1 0 1 0 0 0 1 1 0 1 1 1 0 0 0 0	1296	310
0	−48	0 0 0 0 0 1 1 2 0 1 0 1 0 1 0 0 0 1 1 1 0 1 0 1 0 0 0	648	311
−4	44	0 0 0 0 0 1 1 2 0 1 0 1 1 1 0 0 0 0 1 1 0 0 0 1 0 0 1	648	312
−4	18	0 0 0 0 0 2 0 2 0 0 0 2 0 0 0 2 0 0 0 2 0 2 0 0 0 0 0	36	313
6	−34	0 0 0 0 0 2 0 2 0 0 0 2 0 0 0 2 0 0 1 1 0 1 1 0 0 0 0	324	314
−2	22	0 0 0 0 0 2 0 2 0 0 0 2 0 0 0 2 0 0 2 0 0 0 2 0 0 0 0	108	315
−5	23	0 0 0 0 0 2 0 2 0 0 0 2 1 0 0 1 0 0 1 1 0 0 1 0 1 0 0	648	316
−12	92	0 0 0 0 0 2 0 2 0 0 1 1 1 0 0 1 0 0 0 1 1 1 0 0 1 0 0	162	317
4	−76	0 0 0 0 0 2 0 2 0 0 1 1 1 0 0 1 0 0 1 0 1 0 1 0 1 0 0	648	318
0	128	0 0 0 0 0 2 0 2 0 0 1 1 1 0 0 1 0 0 2 0 0 0 1 0 0 0 1	324	319
−8	40	0 0 0 0 0 2 0 2 0 1 0 1 0 0 0 1 0 1 1 1 0 1 1 0 0 0 0	324	320
−4	108	0 0 0 0 0 2 0 2 0 1 0 1 0 1 0 1 0 0 1 0 1 0 1 0 1 0 0	324	321
4	164	0 0 0 0 0 2 0 2 0 1 0 1 0 1 0 1 0 0 1 1 0 1 0 0 0 0 1	324	322
2	10	0 0 0 0 0 2 0 2 0 1 0 1 1 0 0 0 0 1 1 1 0 0 1 0 1 0 0	648	323
8	−72	0 0 0 0 0 2 1 1 0 0 1 1 0 0 0 1 0 1 1 1 0 1 1 0 0 0 0	1296	324
−6	50	0 0 0 0 0 2 1 1 0 0 1 1 0 1 0 1 0 0 1 0 1 1 0 0 0 1 0	648	325
−2	38	0 0 0 0 0 2 1 1 0 0 1 1 0 1 0 1 0 0 1 1 0 1 0 0 0 0 1	1296	326
5	41	0 0 0 0 0 2 1 1 0 0 1 1 0 1 0 1 0 0 2 0 0 0 1 0 0 0 1	648	327
0	112	0 0 0 0 0 2 1 1 0 0 1 1 1 0 0 0 0 1 1 1 0 0 1 0 1 0 0	1296	328
−6	−30	0 0 0 0 0 2 1 1 0 0 1 1 1 0 0 0 0 1 1 1 0 1 0 0 0 1 0	432	329
−2	158	0 0 0 0 0 2 1 1 0 0 1 1 1 0 0 0 1 0 1 0 1 0 1 0 1 0 0	324	330
−1	−181	0 0 0 0 0 2 1 1 0 0 1 1 1 0 0 0 1 0 2 0 0 0 1 0 0 0 1	648	331
−4	−116	0 0 0 0 0 2 1 1 0 0 1 1 1 0 0 1 0 0 1 1 0 0 1 0 0 0 1	1296	332
2	−54	0 0 0 0 0 2 1 1 0 0 1 1 1 1 0 0 0 0 1 1 0 0 0 0 1 0 1	648	333
−6	34	0 0 0 0 0 2 1 1 0 0 1 1 1 1 0 0 0 0 2 0 0 0 0 0 0 1 1	324	334
4	20	0 0 0 0 0 2 1 1 0 1 1 0 0 1 0 0 0 1 1 1 0 1 0 0 0 0 1	324	335
24	−88	0 0 0 0 1 1 0 1 1 0 1 1 1 0 0 1 0 0 0 1 1 1 0 0 1 0 0	27	336
−8	8	0 0 0 0 1 1 0 1 1 0 1 1 1 0 0 1 0 0 1 0 1 0 1 0 1 0 0	648	337
10	−6	0 0 0 0 1 1 0 1 1 1 0 1 0 0 0 1 0 1 1 1 0 1 1 0 0 0 0	108	338
−8	48	0 0 0 0 1 1 0 1 1 1 0 1 0 0 0 1 1 0 1 0 1 1 1 0 0 0 0	324	339
−2	70	0 0 0 0 1 1 0 1 1 1 0 1 0 0 0 1 1 0 1 1 0 1 0 1 0 0 0	648	340
4	172	0 0 0 0 1 1 0 1 1 1 0 1 0 1 0 1 0 0 1 0 1 1 0 0 0 1 0	324	341
−2	−130	0 0 0 0 1 1 0 1 1 1 0 1 0 1 0 1 0 0 1 1 0 1 0 0 0 0 1	324	342
−12	60	0 0 0 0 1 1 1 0 1 0 1 1 0 0 0 1 1 0 1 0 1 1 1 0 0 0 0	108	343
−4	76	0 0 0 0 1 1 1 0 1 0 1 1 0 0 0 1 1 0 1 1 0 1 0 1 0 0 0	324	344
2	−22	0 0 0 0 1 1 1 0 1 0 1 1 0 0 1 1 0 0 1 1 0 1 0 0 0 1 0	1296	345
4	−108	0 0 0 0 1 1 1 0 1 0 1 1 0 1 0 1 0 0 1 0 1 1 0 0 0 1 0	648	346
0	−136	0 0 0 0 1 1 1 0 1 0 1 1 1 0 0 0 0 1 1 1 0 0 1 0 1 0 0	216	347
6	46	0 0 0 0 1 1 1 0 1 0 1 1 1 0 0 0 1 0 1 0 1 1 0 0 0 1 0	648	348
4	92	0 0 0 0 1 1 1 0 1 0 1 1 1 0 0 0 1 0 1 1 0 0 0 1 1 0 0	648	349
0	−16	0 0 0 0 1 1 1 0 1 0 1 1 1 0 0 0 1 0 1 1 0 1 0 0 0 0 1	1296	350
8	88	0 0 1 0 0 1 1 1 0 0 0 1 0 2 0 1 0 0 1 1 0 1 0 0 0 0 1	216	351
−4	−164	0 0 1 0 0 1 1 1 0 0 0 1 1 1 0 0 1 0 1 1 0 1 0 0 0 0 1	648	352
4	180	0 0 1 0 0 1 1 1 0 0 1 0 1 0 1 0 1 0 2 0 0 0 1 0 0 0 1	324	353
0	144	0 0 1 0 0 1 1 1 0 0 1 0 1 0 1 1 0 0 1 1 0 0 1 0 0 0 1	648	354
−4	236	0 0 1 0 0 1 1 1 0 0 1 0 1 1 0 0 0 1 1 1 0 1 0 0 0 0 1	108	355
0	−128	0 0 1 0 0 1 1 1 0 0 1 0 1 1 0 0 0 1 2 0 0 0 1 0 0 0 1	216	356
−2	−82	0 0 1 0 1 0 1 0 1 0 1 0 1 1 0 0 0 1 2 0 0 0 0 1 0 1 0	324	357
6	6	0 0 1 0 1 0 1 0 1 1 0 0 0 1 1 0 1 0 1 1 0 1 0 0 0 0 1	72	358
2	222	0 0 1 0 1 0 2 0 0 0 2 0 1 0 0 0 0 1 1 0 0 0 0 2 0 1 0	36	359

Table 11. The basis invariants in degree 12: part 5